

## PKU EVENT

Meet Leah and hear  
how she gained  
control of her PKU.

**Palynziq**<sup>®</sup>  
(pegvaliase-pqpz) Injection

How does Leah, a teacher, wife, and mom of 4 control her PKU? Come hear Leah discuss her experience with PALYNZIQ<sup>®</sup> (pegvaliase-pqpz) Injection, a completely different kind of treatment that doesn't require a low-Phe diet!\* You'll learn about PALYNZIQ from a PKU expert and the event is a great way to meet with other adults with PKU.

Please see Important Safety Information, including Boxed Warning for risk of Anaphylaxis, below.



**Atlanta Botanical Garden**  
1345 Piedmont Avenue NE  
Atlanta, GA 30309



**Stephanie Hacker, MS, RD, LDN, CNSC**  
Metabolic Dietitian



**February 29, 2020**  
1:00 PM— 3:00 PM



**Leah Chambless**  
Adult with PKU, taking PALYNZIQ

**Space is limited! Reserve your spot now.**

[www.PKUevents.com/atla2](http://www.PKUevents.com/atla2) or 855-PKU-CALL (855-758-2255)

This special event is intended for adults with PKU (18 years and up) and their family members ages 4 and up.

Limit of 4 family members or caregivers per each person with PKU.

There is no cost to attend this event. It is sponsored by BioMarin Pharmaceutical Inc.

Childcare will not be provided during this event.

Registered attendees will receive tickets to the Atlanta Botanical Garden following the PKU event.

\*The PALYNZIQ Prescribing Information doesn't require a Phe-restricted diet. Talk to your healthcare provider about the treatment plan that's right for you.

What is PALYNZIQ?

PALYNZIQ<sup>®</sup> (Pal-lin-zeek) (pegvaliase-pqpz) Injection is a prescription medication used to lower

## What is PALYNZIQ?

PALYNZIQ® (Pal-lin-zeek) (pegvaliase-pqpz) Injection is a prescription medication used to lower blood levels of Phe (phenylalanine) in adults with PKU (phenylketonuria) who have uncontrolled blood Phe levels above 600 micromol/L (10 mg/dL) on their current treatment. You should discuss the potential benefits and risks of PALYNZIQ with your healthcare provider.

## Important Safety Information

### What is the most important information I should know about PALYNZIQ?

PALYNZIQ can cause a severe allergic reaction (anaphylaxis) that may be life-threatening and can happen any time during treatment with PALYNZIQ.

Severe allergic reactions are a serious but common side effect of PALYNZIQ

- You will receive your first injection of PALYNZIQ in a healthcare setting where you will be closely watched for at least 1 hour after your injection for a severe allergic reaction
- Your healthcare provider will prescribe auto-injectable epinephrine for you, and will teach you (or your caregiver) and your observer, if needed, when and how to use it if you have a severe allergic reaction
- If you have a severe allergic reaction during treatment with PALYNZIQ, you will need to receive an injection of epinephrine immediately and get emergency medical help right away
- Your healthcare provider will decide if you (or your caregiver) are able to give the PALYNZIQ injections, recognize the signs and symptoms of a severe allergic reaction, give an injection of epinephrine, and call for emergency help, if needed
- Your healthcare provider may recommend that an adult observer (or your caregiver) be with you when you give your PALYNZIQ injection and for at least 1 hour after your injection to watch you for signs and symptoms of a severe allergic reaction and, if needed, give you an injection of epinephrine and call for emergency medical help

### Stop injecting PALYNZIQ and get emergency medical care right away if you have any of the following symptoms:

|                                       |  |
|---------------------------------------|--|
| • Fainting (passing out)              | • Swelling of your face, lips, eyes, or tongue |
| • Dizziness or lightheadedness        | • Throat tightness                             |
| • Sudden confusion                    | • Flushed skin                                 |
| • Trouble breathing or wheezing       | • Skin rash, itching, or raised bumps on skin  |
| • Chest discomfort or chest tightness | • Nausea, vomiting, or diarrhea                |
| • Fast heart rate                     | • Losing control of urine or stools            |

**Keep the auto-injectable epinephrine with you at all times during treatment with PALYNZIQ.** Read the Patient Information that comes with the auto-injectable epinephrine that your healthcare provider prescribes for you for more information.

**PALYNZIQ REMS:** PALYNZIQ is available only through a restricted program called the PALYNZIQ REMS (Risk Evaluation and Mitigation Strategy). Talk to your healthcare provider about the PALYNZIQ REMS and how to enroll.

### What should I tell my healthcare provider BEFORE starting PALYNZIQ?

**Tell your healthcare provider about all the medicines you take,** including prescription or over-the-counter medicines, vitamins, and herbal supplements.

**Before injecting PALYNZIQ, talk to your healthcare provider right away** if you cannot or will not use auto-injectable epinephrine to treat a severe allergic reaction.

If you are pregnant or plan to become pregnant while taking PALYNZIQ, talk to your healthcare provider to discuss the risks and benefits of taking PALYNZIQ during pregnancy to you and your unborn baby. If you are breastfeeding or plan to breastfeed, talk to your healthcare provider about the best way to feed your baby if you take PALYNZIQ.

**Before injecting PALYNZIQ, read the Medication Guide and Instructions for Use** that come with your PALYNZIQ injection.

### What should I watch for AFTER starting PALYNZIQ?

**PALYNZIQ may cause serious side effects, including:**

- Severe allergic reactions (anaphylaxis)
- Other allergic reactions to PALYNZIQ can happen during treatment with PALYNZIQ. Contact your healthcare provider right away if you have any of the following symptoms of an allergic reaction including: rash, itching, or swelling of the face, lips, eyes, or tongue. Your healthcare provider may change your dose of PALYNZIQ, stop your treatment with PALYNZIQ for a period of time, or prescribe medicine for you to take before your PALYNZIQ injection to help reduce the symptoms of an allergic reaction

**The most common side effects of PALYNZIQ** include injection site reactions (such as redness, itching, pain, bruising, rash, swelling, or tenderness), joint pain, headache, skin reactions that spread and last at least 14 days (such as itching, rash, or redness), itching, nausea, stomach pain, mouth and throat pain, vomiting, cough, diarrhea, feeling very tired, and low levels of Phe in your blood.

**These are not all of the possible side effects of PALYNZIQ. Speak with your healthcare provider right away about any side effects.**

### Important notes

Blood Phe testing and diet

- Your healthcare provider will monitor your blood Phe concentrations during PALYNZIQ treatment
- Monitor the amount of protein and Phe that you eat or drink. Your healthcare provider may change the amount of protein and Phe you should have in your diet during treatment with PALYNZIQ, depending on the levels of Phe in your blood. Follow your healthcare provider's instructions about the amount of protein and Phe you should have in your diet

### Missed dose